


# Ramachandran Plot

## pdb2ibz




### Plot statistics

Residues in most favoured regions [A,B,L]	1672	87.7%
Residues in additional allowed regions [a,b,l,p]	223	11.7%
Residues in generously allowed regions [-a,-b,-l,-p]	9	0.5%
Residues in disallowed regions	2	0.1%
-----		
Number of non-glycine and non-proline residues	1906	100.0%
Number of end-residues (excl. Gly and Pro)	21	
Number of glycine residues (shown as triangles)	141	
Number of proline residues	99	
-----		
Total number of residues	2167	

Based on an analysis of 118 structures of resolution of at least 2.0 Angstroms and R-factor no greater than 20%, a good quality model would be expected to have over 90% in the most favoured regions.

# Ramachandran plots for all residue types


## pdb2ibz



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

# Ramachandran plots for all residue types


## pdb2ibz



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

# Ramachandran plots for all residue types


## pdb2ibz



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

# Chi1-Chi2 plots


## pdb2ibz



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

# Chi1-Chi2 plots







## pdb2ibz



Numbers of residues are shown in brackets. Those in unfavourable conformations (score < -3.00) are labelled. Shading shows favourable conformations as obtained from an analysis of 163 structures at resolution 2.0A or better.

# Main-chain parameters

pdb2ibz








Plot statistics

Stereochemical parameter	No. of data pts	Parameter value	Comparison values		No. of band widths from mean	
			Typical value	Band width		
a. %-tage residues in A, B, L	1906	87.7	79.8	10.0	0.8	Inside
b. Omega angle st dev	2151	1.3	6.0	3.0	-1.6	BETTER
c. Bad contacts / 100 residues	5	0.2	7.6	10.0	-0.7	Inside
d. Zeta angle st dev	2026	1.4	3.1	1.6	-1.0	BETTER
e. H-bond energy st dev	1394	0.7	0.9	0.2	-0.8	Inside
f. Overall G-factor	2167	0.3	-0.5	0.3	2.6	BETTER

# Side-chain parameters

## pdb2ibz



pdb2ibz


### Plot statistics

Stereochemical parameter	No. of data pts	Parameter value	Comparison values		No. of band widths from mean	
			Typical value	Band width		
a. Chi-1 gauche minus st dev	257	12.5	20.9	6.5	-1.3	BETTER
b. Chi-1 trans st dev	608	14.6	21.2	5.3	-1.2	BETTER
c. Chi-1 gauche plus st dev	892	13.8	19.8	4.9	-1.2	BETTER
d. Chi-1 pooled st dev	1757	14.0	20.5	4.8	-1.3	BETTER
e. Chi-2 trans st dev	495	14.5	22.0	5.0	-1.5	BETTER




# Residue properties pdb2ibz


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion



c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb2ibz


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb2ibz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

### d. Secondary structure & estimated accessibility



Key:- Helix Beta strand Random coil Accessibility shading: Buried Accessible


### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb2ibz


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion




Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb2ibz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb2ibz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb2ibz


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb2ibz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors






# Residue properties pdb2ibz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion



### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb2ibz


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




Key:- Helix Beta strand Random coil Accessibility shading: Buried Accessible


e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors




c = cis-peptide

# Residue properties pdb2ibz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb2ibz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion



### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb2ibz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb2ibz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb2ibz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb2ibz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors






# Residue properties pdb2ibz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb2ibz


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion



c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb2ibz


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb2ibz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion



### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb2ibz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors




# Residue properties pdb2ibz


a. Absolute deviation from mean Chi-1 value (excl. Pro)



b. Absolute deviation from mean of omega torsion




c. C-alpha chirality: abs. deviation of zeta torsion




Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal

d. Secondary structure & estimated accessibility




e. Sequence & Ramachandran regions Most favoured Allowed Generous Disallowed



f. Max. deviation (see listing)




g. G-factors




# Residue properties pdb2ibz


### a. Absolute deviation from mean Chi-1 value (excl. Pro)



### b. Absolute deviation from mean of omega torsion




### c. C-alpha chirality: abs. deviation of zeta torsion



Highlighted residues are those that deviate by more than 2.0 st. devs. from ideal


### d. Secondary structure & estimated accessibility




### e. Sequence & Ramachandran regions



### f. Max. deviation (see listing)




### g. G-factors



c = cis-peptide

# Main-chain bond lengths

## pdb2ibz




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.



# Main-chain bond angles

## pdb2ibz




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

# Main-chain bond angles

## pdb2ibz




Black bars > 2.0 st. devs. from mean.

Solid and dashed lines represent the mean and standard deviation values as per Engh & Huber small-molecule data.

# RMS distances from planarity

## pdb2ibz




Histograms showing RMS distances of planar atoms from best-fit plane.  
 Black bars indicate large deviations from planarity: RMS dist > 0.03 for rings, and > 0.02 otherwise.

▶ signifies data points off the graph in the direction shown.


# Distorted geometry pdb2ibz

## Main-chain bond angles



Bond angles differing by > 10.0 degrees from small-molec values. Values shown: "ideal", actual, diff.


## Planar groups



# Distorted geometry

## pdb2ibz

### Planar groups (contd)



Sidechains with RMS dist. from planarity > 0.03A for rings, or > 0.02A otherwise. Value shown is RMS dist.